arXiv:1405.7687v2 [physics.gen-ph] 9 Jun 2015

Pre-big bang scenario and the WZW model

Marcel Jacon

June 10, 2015

Abstract

Extensive studies of pre-big bang scenarios for Bianchi-I type universe have been made, at various approximation levels. Knowing the solution of the equations for the post-big bang universe, the symmetries of the equations ("time reversal and scale dual transformations") allow the study of pre-big bang solutions. However, they exhibit singularitites in both the curvature and the dilaton kinetic energy.

Calculating the β equations for the Non-Linear Sigma model, at the first loop approximation and imposing conformal invariance at this level, lead to equations of motion that simply state that the curvature must be nil , which in turn allows the utilization of groups to solve the β equations. This is what is done in the Weiss-Zumino-Witten (WZW) model.

In this article, we will show that using the WZW model on SU_2 ,some of the difficulties encountered in the determination of the pre and post big-bang solutions are eliminated, leading to realistic solutions for the evolution of the universe and giving an explanation to the actually observed acceleration of the expansion.

1 Introduction:

Extensive studies of pre-big bang scenarios for Bianchi-I type universe have been made at various approximation levels [1,2] and references herein. The gravitational (massless, bosonic) sector of the string action contains not only the metric, but also ,at least one more fondamental field, the dilation ϕ.The corresponding tree-level action lead to cosmological equations which have been established in the case where the (NS-NS) two form $B_{\mu\nu}$=0, but including the contribution of perfect fluid sources.These equations are invarariant under "time reversal transformation, but also under "scale dual transformations". Knowing the solution of the equations for the post-big bang universe, these symmetry allow the study of pre-big bang solutions. The exact integration of the string cosmology equations in the fully anisotropic case can be performed but lead to solutions which exhibit singularitites in both the curvature and the dilaton kinetic energy. The solutions associated with the pre and post-big bang branches, being disconnected by a singularity, are not appropriate to describe the whole transition between the two regimes. Particular examples of regular solutions

may be obtained, but one may also expect that the regularisation of the big bang singularity need also to introduce the effects of higher order loop and α' corrections.

The Polyakov action may be modified to incorporate the effect of massless excitation, and leads to theNon-Linear Sigma model [3,4]. Calculating the β equations for the Non-Linear Sigma (NLS) model at the first loop approximation and imposing conformal invariance at this level, leads to equations of motion that simply state that the curvature must be nil, which in turn allows the utilization of group manifolds to solve the β equations. This is what is done in the Weiss-Zumino-Witten (WZW) model [5]

In this article, we will show how, using the WZW model on SU_2 ,some of the difficulties encountered in the determination of the pre and post big-bang solutions are eliminated, leading to realiistic solutions for the evolution of the universe.

2 The WZW-model on SU_2.

An element of SU_2 is parametrized by means of the well-known Euler angles (α, β, γ):

$$g = \exp(\alpha\tau_3)\exp(\beta\tau_2)\exp(\gamma\tau_3)$$

where the τ_i matrices are given in terms of Pauli matrices $\tau_k = \frac{\sigma_k}{2i}$, and the Euler angles have values in the ranges $0 \leq \alpha < 2\pi$, $0 \leq \beta < \pi$, $0 \leq \gamma < 2\pi$

Using the expressions for the Maurer-Cartan forms [6] , one deduces the line element:

$ds^2 = d\alpha^2 + d\beta^2 + d\gamma^2 + 2d\alpha d\gamma \cos\beta$ which determines the g matrix

$$g = \begin{pmatrix} 1 & 0 & \cos\beta \\ 0 & 1 & 0 \\ \cos\beta & 0 & 1 \end{pmatrix} \tag{2.0.1}$$

whose invere is

$$g^{-1} = \frac{1}{\sin^2\beta} \begin{pmatrix} 1 & 0 & -\cos\beta \\ 0 & \sin^2\beta & 0 \\ -\cos\beta & 0 & 1 \end{pmatrix} \tag{2.0.2}$$

From the elements of these matrix, one easily obtain the Christoffel symbols and finally the Ricci tensors elements and the curvature scalar

$$R = R_{\mu\nu} g^{\mu\nu} = \frac{3}{2} \tag{2.0.3}$$

A space of constant curvature is a conformally flat space, so that its Weyl tensor vanishes identically.

2.1 The evolution of the universe:

When studying the evolution on the universe, we work in the *S-frame representation* of the string effective action[2]. Assuming that $B_{\mu\nu}$ is vanishing and that the matter sources can be represented by a fluid, in the synchronous gauge of the comoving frame, we have:

$$\tilde{g}_{\mu\nu} = diag(1,\text{-}\tilde{a}^2\delta_{ij}),\ \tilde{a} = \tilde{a}(t),\ \tilde{\phi} = \tilde{\phi}(t)$$

$$\tilde{T}^{\nu}_{\mu}\text{=diag}(\tilde{\rho} - \tilde{p}\delta_{ij}),\ \tilde{\rho} = \tilde{\rho}(t),\ \tilde{p} = \tilde{p}(t),\ \tilde{\sigma} = \tilde{\sigma}(t)$$

These definitions and the use of the cosmic time as the time coordinate leads to the well-known Roberson-Walker metric

$$ds^2\text{=}dt^2 - \tilde{a}^2(t)\left[(dr^2/1 - Kr^2) + r^2(d\theta^2 + sin^2\vartheta d\phi^2)\right]$$

In this formula, K has dimension L^{-2} and $\tilde{a}$(t) is dimensionless.

In order to preserves the SU_2 character of the 3-dimensional spatial subgroup and keep its interesting symmetry properties, such as a constant curvature (i.e. independent of the spatial coordinates), for a fixed t, we will work in the Euler Angles frame which is described now.

We look for a coordinate system (ξ, η, ζ) which diagonalizes the matrix diag (-a^2,-a^2,-a^2)×g

(2.1.1)

where a depends on t .

One has the following transformation:

$$\begin{pmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{-1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{pmatrix} : \begin{pmatrix} -a^2 & 0 & -a^2\cos\beta \\ 0 & -a^2 & 0 \\ -a^2\cos\beta & 0 & -a^2 \end{pmatrix} \begin{pmatrix} \frac{1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \\ 0 & 1 & 0 \\ \frac{-1}{\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{pmatrix}$$

$$= \begin{pmatrix} -a^2(1-\cos\beta) & 0 & 0 \\ 0 & -a^2 & 0 \\ 0 & 0 & -a^2(1+\cos\beta) \end{pmatrix}$$

From which it follows:

$$-a^2(d\alpha^2 + d\beta^2 + d\gamma^2 + 2\cos\beta d\alpha d\gamma) = -a^2((1-\cos\beta)d\xi^2 + d\eta^2 + (1+\cos\beta)d\zeta^2)$$

$$\text{with } \left[d\xi = (d\alpha - d\gamma)/\sqrt{2}, d\eta = d\beta, d\zeta = (d\alpha + d\gamma)/\sqrt{2}\right]$$

The 4-dimensional line element is

$$\left[ds^2 = dt^2 - a^2((1-\cos\beta)d\xi^2 + d\eta^2 + (1+\cos\beta)d\zeta^2)\right]$$

from which one sees that the scale factor a has the dimension L

In the S-frame, fixing $r = r_0$, an arbitrary value, gives $ds^2 = dt^2 - \tilde{a}^2(t)\left[r_0^2(d\theta^2 + \sin^2\theta d\phi^2)\right]$.

On the other hand, defining $d\theta = d\eta$ and $\sin^2\theta d\phi^2 = 2(\sin^2(\beta/2)d\xi^2 + \cos^2(\beta/2)d\zeta^2)$ gives the expected result $a = \tilde{a}(t)\times r_0$.

(2.1.2)

In this coordinate system, we will see that the curvature scalar is given by the following constant, generalizing (2.0.3):

$$R\text{=-}(3/2a^2)$$

(2.1.3)

As a consequence, the β-equations for a fixed t at the first loop in string perturbation theory, imposing that the theory is conformal invariant at this order, are satisfied .

(2.1.4)

We study the graviton-dilaton system , setting $B_{\mu\nu} = 0$, but including perfect fluid sources. Therefore ϕ =ϕ(t) ,T^{ν}_{μ} =diag $(\rho, -p_i\delta^j_i)$, ρ=$\rho(t)$,p_i= $p_i(t)$, σ =$\sigma(t)$, where

$T_{\mu\nu}$represente the tenseur current density of the matter sources andσ the scalar charge density.

We first calculate the components of the Christoffell connection Γ, and deduce the components of the Ricci tensor:

From

$$\Gamma^k_{ij}=\frac{1}{2}g^{kl}(\partial_i g_{jl}+\partial_j g_{il}\text{-}\partial_l g_{ij})$$

we deduce

$$\Gamma^i_{0i} = \frac{\dot{a_i}}{a_i} = H_i;\ \Gamma^0_{ii} = \dot{a}_i a_i;\ \Gamma^1_{12} = \Gamma^1_{21} = \frac{\partial_2 a_1}{a_1} = \frac{1}{2}\frac{a_3}{a_1};\ \Gamma^3_{23} = \Gamma^3_{32}=\frac{\partial_2 a_3}{a_3} = -\frac{1}{2}\frac{a_1}{a_3}:$$
$$\Gamma^2_{11} = -\frac{a_1}{a_2^2}\partial_2 a_1 = -\frac{1}{2}\frac{a_1 a_3}{a_2^2};\ \Gamma^2_{33} = -\frac{a_3}{a_2^2}\partial_2 a_3 = \frac{a_3 a_1}{a_2^2}$$

One obtains the following expressions, which apart from a constant, are similar to those calculated by M. Gasperini in the *S-frame* (see Eqns. 4.10 of [2])

$$R^0_0\text{=-}\textstyle\sum_i(\dot{H}_i+H_i^2),$$

$$R^i_i\text{=-}\dot{H}_i\text{-}H_i(\textstyle\sum_k H_k)\text{-}1/2a^2$$

(2.1.5)

The associated scalar curvature is:

$$R\text{=-}\textstyle\sum_i(2\dot{H}_i+H_i^2)\text{-}(\sum_i H_i)^2\text{-}3/2a^2$$

showing the above result (Eqn 2.1.3) for the three dimensional spatial manifold.

(2.1.6)

We retain also the following equations for the dilaton field:

$$(\nabla\phi)^2=\dot{\phi}^2,\ \nabla^2\phi=\ddot{\phi}+\dot{\phi}\sum_i H_i,\ \nabla_0\nabla^0\phi=\ddot{\phi},\ \nabla_i\nabla^j\phi=\dot{\phi}H_i\delta_i{}^j$$

We have also the following relations between the quantities in the E and S frames:

$$H_1=H_2=H_3=\frac{\dot{a}}{a}=\frac{\dot{\tilde{a}}}{\tilde{a}}=\tilde{H}$$

So the equations for the theory are the same, in the S frame or in the E frame, except the remaining constant in theR_i^i .

it follows that we can use the results already obtained by M. Gasperini in the S-frame.

(2.1.7)

Setting $2{\lambda_s}^{d-1}=1$,we have the Euler-Lagrange equation for the dilaton equation (analogous of Eqn 2.24 of [2].

$$\ddot{\phi}\text{-}\dot{\phi}^2+2\dot{\phi}\textstyle\sum_i H_i\text{-}\sum_i(2\dot{H}_i+H_i^2)\text{-}(\sum_i H_i)^2\text{-}3/2\mathrm{a}^2+V\text{-}\frac{\partial V}{\partial\phi}=\frac{1}{2}e^{\phi}\sigma$$

(2.1.8)

The $(0,0)$ component of (Eqn.2.24) in [2] gives

$$\dot{\phi^2}\text{-}2\dot{\phi}\textstyle\sum_i H_i+(\sum_i H_i)^2\text{-}\sum_i(H_i^2)\ +(3/2a^2)\text{-}V=e^{\phi}\varrho$$

(2.1.9)

While the diagonal part of the space component gives:

$$\dot{H}_i\text{-}H_i(\dot{\phi}\text{-}\textstyle\sum_k H_k)+\frac{1}{2}\frac{\partial V}{\partial\phi}\text{-}(1/4a^2)=\frac{1}{2}e^{\phi}(p_i-\frac{\sigma}{2})$$

(2.1.10)

These equations are simplified, introducing the "shifted dilaton variable " $\bar{\phi}$

$$\bar{\phi}=\phi\text{-}\ln(a^3),\dot{\bar{\phi}}=\dot{\phi}\text{-}\textstyle\sum_i H_i$$

(2.1.11)

and the shifted variables for the fluid:

$$\bar{\varrho}=\rho(a^3),\ \bar{p}=p(a^3),\bar{\sigma}=\sigma(a^3)$$

then, we obtain the analogous of Eqns 4.39,4.40,4.41 in [2]:

$$\dot{\bar{\phi}}^2\text{-}\textstyle\sum_i(H_i^2)\text{-}\frac{3}{2a^2}\text{-}V=\bar{\rho}e^{\bar{\phi}}$$

$$\dot{H}_i\text{-}H_i\dot{\bar{\phi}}+\frac{1}{2}\frac{\partial V}{\partial\bar{\phi}}\text{-}\frac{1}{4a^2}=\frac{1}{2}e^{\bar{\phi}}(\bar{p}_i\text{-}\frac{\bar{\sigma}}{2})$$

$$2\ddot{\bar{\phi}}\text{-}\dot{\bar{\phi}}^2\text{-}\sum_i(H_i^2)\text{-}\frac{3}{2a^2}+V\text{-}\frac{\partial V}{\partial\bar{\phi}}=\frac{1}{2}\bar{\sigma}e^{\bar{\phi}} \tag{2.1.12}$$

From these equations, we deduce the following conservation equation:

$$\dot{\bar{\rho}}+\sum_i H_i p_i=\frac{1}{2}\sigma(\dot{\bar{\phi}}+\sum_i H_i)+e^{-\bar{\phi}}\left[\frac{\partial V}{\partial\bar{\phi}}\sum_i H_i+\frac{3}{a^2}\bar{\phi}-\frac{1}{2a^2}\sum_i H_i-\frac{3}{2}\frac{d}{dt}\frac{1}{a^2}\right]$$

However V is not a scalar under general coordinates transformations, and it is impossible to define a potential which can be directly inserted as a scalar into the covariant action. However, it has been shown that the action and the corresponding equations of motion can be written in a generalized form which is invariant under general coordinates transformations using for the potential a non-local variable.

.The result of the calculations is that the second Eqn. (2.1.12) is replaced by the simpler one (2.1.13) [2,7]:

$$\dot{H}_i\text{-}H_i\dot{\bar{\phi}}\text{-}\frac{1}{4a^2}=\frac{1}{2}e^{\bar{\phi}}\bar{p}_i \tag{2.1.13}$$

leading to the modified conservation equation:

$$\dot{\bar{\rho}}+\sum_i H_i p_i=\frac{1}{2}\sigma\dot{\bar{\phi}}+e^{-\bar{\phi}}\left[\dot{\bar{\phi}}\frac{3}{2a^2}-\frac{1}{2a^2}\sum_i H_i-\frac{3}{2}\frac{d}{dt}\frac{1}{a^2}\right]$$

as well as the Eqn for the Dilaton field $\phi(\bar{t})$: (Eqn 4A.27 of ref.2)

$$\ddot{\bar{\phi}}+\sum_i H_i\dot{\bar{\phi}}\text{-}\dot{\bar{\phi}}^2+2\lambda_s^{d-1}(V-\frac{1}{2}\frac{\partial V}{\partial\bar{\phi}})+\lambda_s^{d-1}e^{\phi}(\rho-\sum_i p_i)+\frac{1}{2}(d-1)\lambda_s^{d-1}e^{\phi}\sigma=0 \tag{2.1.14}$$

Using these equations, we can now study models for the evolution of the Universe.

2.2 Models for the evolution of the Universe:

Instead of introducing a time parameter x as in [2] and get a full analytical result, we calculate a numerical solution of the differential equations of motion [**?**]. However the relations between both methods are easily found. It has been shown by M. Gasperini and Veneziano [2,7] that the cosmological equations are rigorously solved, in the isotropic case and for the vacuum ($T_{\mu\nu}=0=\sigma$) by the solutions

$$A(t)=A_0\left[\frac{t}{t_0}+\left(1+\frac{t^2}{t_0^2}\right)^{1/2}\right]^{1/\sqrt{d}} \tag{2.2.1}$$

and

$$\bar{\phi} = -\frac{1}{2}\ln\left[\sqrt{V_0}t_0\left(1+\frac{t^2}{t_0^2}\right)\right]$$

Our equations (**??**) are slightly different with the presence of the factors $\frac{1}{a^{2,}}$, leading to the following conditions, which must relate the scale factor to the sources properties:

$$\frac{1}{a^2} = -\frac{1}{3}e^{\bar{\phi}}\bar{\sigma} = -2e^{\bar{\phi}}\bar{p} = -\frac{2}{3}e^{\bar{\phi}}\bar{\rho} \Longrightarrow \bar{\rho} = 3\bar{p},\ \bar{\sigma} = 2\bar{\rho} \tag{2.2.2}$$

This shows that the solutions (2.2.2) are even valid in the presence of matter, provided the conditions [2] $\gamma = 1/3$, $\gamma_0 = 2$ are satisfied, i.e. in the case of a pure radiation field :this is an improvement over previous results.

The curves representing $H(t)$ and $\bar{\phi}$ are similar to those of [2] , (Fig. 4.7) with a bell-like shape for the curvature and the dilaton kinetic energy. They show a pre-big bang inflationary evolution, followed by a decelerated expansion.

These solutions exhibit no possible acceleration of the universe expansion, which is actually observed. This demonstrates that such an acceleration is due to the presence of another kind of field, such as dark energy.

The most recent experimental results on the observation of an acceleration of the actual expansion of the universe is found in [**?**]. From the results published in this report, we can perform a fitting of the continuous curve of $\dot{\tilde{A}}(t)$ represented on Fig.21 of [**?**]. For that, we must first numerically solve the first order differential equation which determine $\tilde{A}(t)$:

$$\dot{\tilde{A}}(t) = \tilde{A}(t) \times H_0(\Omega_\Lambda + \Omega_M(1+z)^{\dot{3}} + (1-\Omega_\Lambda-\Omega_M)(1+z^2))^{1/2} \tag{2.2.3}$$

Performing this operation is not easy, because this is a stiff equation which must be solved by efficient algorithms. This is done using the Mathematica package [**?**]. Concerning the variables, we will use either t or the redshift parameter z , which taking arbitrarily $\tilde{A(0)} = 1$, are related by

$$1 + z(t) = \frac{1}{y(t)},\ y(t) = \tilde{A}(t) \tag{2.2.4}$$

A numerical fitting of $z(t)$ Taking arbitrarily $\tilde{A}(0) = 1$, is:

$$\begin{aligned} 1/\tilde{A}(t) \quad = \quad & 1.0 - 70.62831490791527t + 3740.638505154311t^2 \\ & -79103.96903009895t^3 - 1.578335812532445 \times 10^7 t^4 \\ & -6.744184076495469` \times 10^9 t^5 + 8.136104416644515` \times 10^{11} t^6 \\ & +1.1127413276575105 \times 10^{14} t^7 - 5.506299532617992 \times 10^{15} t^8 \\ & -8.075146184886697 \times 10^{17} t^9 \end{aligned}$$

(2.2.5)

valid in the interval $-0.011 \leqslant t \leqq 0.008$, $t = 0$ being the present epoch. In fact the unit for t is very large: relatively to the present epoch, $\triangle z = 0.5$ corresponds to $\triangle t = 0.0066$s. If we consider that the expansion of the universe began to accelerate at $\triangle T = 5My$ (cosmic time)[**?**], it follows $\triangle T/\triangle t = 2.38 \times 10^{19}$. So the time interval between $t_1 = -0.011$ and $t_2 = 0.008$ is $\triangle T = 14.35My$ and covers the whole life of the universe.

$\tilde{A}(t)$ is an increasing function of t. The curve $\dot{\tilde{A}}(t)$, when expressed in terms of the redshift parameter $z(t)$, reproduces the continuous curve of Fig. 20 [**?**].

The equation which determines $\bar{\phi}(t)$ is deduced from (**??**):

$$\ddot{\bar{\phi}} + \left(1 - \frac{\gamma_0}{2}\right)\frac{1}{\gamma}H\dot{\bar{\phi}} - \frac{\partial V}{\partial \bar{\phi}} = dH^2 + \left(1 - \frac{\gamma_0}{2}\right)\frac{1}{\gamma}\dot{H} + \frac{1}{2a^2}\left[3 - \frac{1}{2\gamma}\left(1 - \frac{\gamma_0}{2}\right)\right] \tag{2.2.6}$$

Now, the solutions that we have adopted for $A(t)$, $\bar{\phi}(t)$, $V(\bar{\phi})$ are not rigorous solutions of (**??**). To get a rigorous solution, we must add a correction δV to the potential to compensate for the other components of the cosmic field, such that

$$\begin{aligned} -\frac{3}{2a^2} - \delta V &= e^{\bar{\phi}}(\bar{\rho} - \bar{\rho}_r) \\ -\frac{1}{4a^2} = \frac{1}{2}e^{\bar{\phi}}(\bar{p} - \bar{p}_r) &= \frac{1}{2}e^{\bar{\phi}}\gamma(\bar{\rho} - \bar{\rho}_r) \\ -\frac{3}{2a^2} + \delta V - \frac{\partial}{\partial \bar{\phi}}(\delta V) &= \frac{1}{2}e^{\bar{\phi}}(\bar{\sigma} - \bar{\sigma}_r) \end{aligned} \tag{2.2.7}$$

From which results:

$$\begin{aligned} -\frac{1}{a^2} &= 2e^{\bar{\phi}}\gamma(\bar{\rho} - \bar{\rho}_r) \\ \delta V &= e^{\bar{\phi}}(3\gamma - 1)(\bar{\rho} - \bar{\rho}_r) \\ \frac{\partial}{\partial \bar{\phi}}(\delta V) &= e^{\bar{\phi}}\left(6\gamma - 1 - \frac{\gamma_0}{2}\right)(\bar{\rho} - \bar{\rho}_r) \end{aligned} \tag{2.2.8}$$

It follows that Eqn (**??**) is now

$$\ddot{\bar{\phi}} + \left(1 - \frac{\gamma_0}{2}\right)\frac{1}{\gamma}H\dot{\bar{\phi}} = dH^2 + \left(1 - \frac{\gamma_0}{2}\right)\frac{1}{\gamma}\dot{H} - e^{\bar{\phi}}\frac{\gamma_0}{2}(\bar{\rho} - \bar{\rho}_r) = dH^2 + \left(1 - \frac{\gamma_0}{2}\right)\frac{1}{\gamma}\dot{H} + \frac{1}{4a^2}\frac{\gamma_0}{\gamma}$$

with

Figure 1: The density ρ as a function of z

$$\gamma_0 = 2, \gamma = \frac{1}{3} \tag{2.2.9}$$

With these values for the two constant, Eqn (**??**) takes the remarkable simple form

$$\ddot{\bar{\phi}} = dH^2 + \frac{1}{4a^2}(\gamma_0/\gamma) \tag{2.2.10}$$

We have the following values for the density [**?**]:

$$\rho = \Omega_M(1+z)^3 + \Omega_\Lambda + (1 - \Omega_\Lambda - \Omega_M)(1+z)^2$$

A plot of the density as a function of z is given in Fig. 2.

We make a numerical integration of Eqn. (**??**), we need to evaluate $H(t)$ and $\dot{H}(t)$ which are given by the following expressions:

$$H(t) = \frac{\dot{a}(t)}{a(t)},\ \dot{H}(t) = \left(\frac{\ddot{a}(t)}{a(t)}\right) - \left(\frac{\dot{a}(t)}{a(t)}\right)^2$$

$1/\tilde{A}(t)$ being given by (**??**).

We have also:

$$H(z) = 69.9868 + 28.7795 \times z + 22.612 \times z^2 - 2.45536 \times z^3$$

We obtain the curve given in Fig. 2.2, showing the evolution of $\bar{\phi}(t)$ with the initial condition $\bar{\phi}(t = -0.0066\text{s}) = 0$, fixing the beginning of an accelerating expansion at $T = -5My$. $\bar{\phi}(t)$, correlating to the evolution of the expansion, is decreasing during the decelerating expansion, and then increasing during the accelerating expansion phase.

Matching of the solutions: To get a full description of the universe evolution, we must first propagate Gasperini solution as soon as the radiation field dominates,- as we have seen that the bell-shaped GV-solution is still valid. For later time, we must use the ΛCDM solution , matching both at the junction point.

This is done in the following way.

The GV-solution (Gasperini-Veneziano) is



Figure 2: The dilaton $\bar{\phi}(t)$



Figure 3: The global solution for the dilaton field $\bar{\phi}(t)$.

$$H(t) = \left(d \times (t^2 + t_1^2)\right)^{-0.5},\ \dot{\bar{\phi}}(t) = -t/(t^2 + t_1^2)$$

Matching $H(\Lambda CDM)$ and H(GV) at $t = -0.006$ fixes $t_1 = 0.000678617$. It results that $\dot{\bar{\phi}}(t = -0.006) = 166.667$. Taking $\bar{\phi}(-0.006) = 0$ we get the curve given in Fig. 3, showing the increasing of $\bar{\phi}(t)$.

3 Conclusion.

In this article, I have developped models for the evolution of the universe, using the Weiss-Zumino-Witten method on the SU_2 group. Evolution equations for the fields variables have been established in the simplest case where the graviton-dilaton system is described by the dilaton field $\phi(t)$. Solution are numerically computed, taking care of the stiffness of the equations . These solutions describe well the actual acceleration of the expansion which, according to the most present measurements began around 5 billion years ago. So this simple models show the ability of string theory, to describe the actually observed evolution of the universe, giving an interpretation in terms of the dilaton field.

References

[1] M.Gasperini and G. Veneziano, String Theory and Pre-Big bang Cosmology. Contribution to the book: Beyond the Big Bang, edited by Ruediger Vaas (Frontier Collection Series, Springer Verlag, Heidelberg, 2007)

[2] M. Gasperini, Elements of String Cosmology (Cambridge University Press, Cambridge 2007)

[3] E.S. Fradkin , A.A. Tseytlin: Quantum String Theory effective action , Nucl. Phys. B **261**, 1 (1985)

[4] C.G. Callen, D. Freedan, E.I.Martinec, M.J. Perry: String in background fields, Nucl. Phys. B **262**, 593 (1985)

[5] E. Witten: Non-Abelian bosonization in two dimensions, Comm. Math. Phys. **93**, 455 (1984)

[6] Exercises on Differential Geometry of Lie Groups; http://compalg-jinr.ru/CAGroup/Palii/GeomSU23.pdf

[7] M. Gasperini, Dilaton Cosmology and Phenomenology, hep-th/0702166v2 (2007)

[8] Wolfrarm Mathematica 9, Home Edition

[9] La Recherche, Septembre 2013 : Energie noire: pourquoi l'Univers accélère.

[10] N.G. Busca et al., arXiv:1211.2616 (2012)